

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets

(11) Publication number:

0 036 636**B1**

(12)

EUROPEAN PATENT SPECIFICATION(45) Date of publication of patent specification: **15.02.84**(51) Int. Cl.³: **C 07 C 107/06,**
C 09 B 29/14(21) Application number: **81102068.4**(22) Date of filing: **19.03.81**(54) **A method and intermediates for preparing 3,3'-azo-bis-(6-hydroxy benzoic acid).**(30) Priority: **26.03.80 SE 8002321**(43) Date of publication of application:
30.09.81 Bulletin 81/39(45) Publication of the grant of the patent:
15.02.84 Bulletin 84/7(84) Designated Contracting States:
AT BE CH DE FR GB IT LI LU NL SE(56) References cited:
DE - C - 278 613
DE - C - 590 321
FR - A - 754 674
FR - A - 2 161 104
GB - A - 408 676
GB - A - 1 167 176
US - A - 4 045 429(73) Proprietor: **PHARMACIA AKTIEBOLAG**
Rapsgatan 7 Box 604
S-751 25 Uppsala (SE)(72) Inventor: **Agback, Karl Huberg**
Mimervägen 7
S-754 40 Uppsala (SE)
Inventor: **Nygren, Alf Sigurd**
Mänstavägen 58
S-740 60 Örbyhus (SE)(74) Representative: **Kraus, Walter, Dr. et al,**
Patentanwälte Dres. Kraus & Welsert
Irmgardstrasse 15
D-8000 München 71 (DE)

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European patent convention).


Courier Press, Leamington Spa, England.

EP 0 036 636 B1

0 036 636

A method and intermediates for preparing 3,3'-azo-bis-(6-hydroxy benzoic acid)

The present invention relates to a new method and new intermediates for preparing 3,3'-azo-bis(6-hydroxy benzoic acid) of the formula



and salts thereof. The new method according to the invention makes it possible to prepare the compound of formula I and its salts, especially pharmaceutically acceptable salts, in very high purity, for example 98% and more, and with good yields.

The compound of formula I known as such and has found commercial use as a dyestuff. A known method for preparing the compound I involves alkaline hydrolysis of 3,3'-azo-6-chloro-6'-hydroxy-dibenzoic acid at high temperature, possibly in the presence of copper: see for example DE Patent 278613. In an alternative known method of preparation the compound in question is prepared by a corresponding alkaline hydrolysis of 3,3'-azo-bis-(6-chlorobenzoic acid): see for example GB Patent 408676. In a third known method of preparation 5-amino salicylic acid is diazotized and coupled with salicylic acid in the presence of trivalent chromium, whereupon the chromium complex formed is decomposed for setting free the compound I: see for example DE Patent 590321.


The first mentioned method of preparation is the one which is used commercially. The product obtained in this hydrolysis of 3,3'-azo-6-chloro-6'-hydroxy-dibenzoic acid is not pure, but contains comparatively great amounts of starting material (e.g. 20 to 30%). Further, the desired end product I is not stable under the severe hydrolysis conditions but is decomposed, i.e. to 5-amino salicylic acid.

It would theoretically be possible to separate the end product I from the starting material, for example by using liquid chromatographic analysis technique. Such preparation method are, however, economically impossible to apply for practical preparative use. Because of the very similar properties of the starting and the end compounds it has not either been possible to separate them by crystallisation.

The alternative synthesis route via the chromium complex of the compound I does not either yield any pure product since impurities of chromium always are present. Because of such impurities the compound I, when prepared in this manner, cannot be used as e.g. a drug.

It is an object of the present invention to provide a technically simple and economically advantageous new method of preparing 3,3'-azo-(bis-6-hydroxy-benzoic acid) of formula I, in which the desired product is obtained in considerably higher purity than in the known methods of preparation. In particular, the invention makes it possible to prepare the compound I in a purity of at least 98%, thereby making it possible to use the compound in question as a drug, for example for treating inflammatory intestinal diseases such as, for example, ulcerous colitis. This use is described in our simultaneously herewith filed EPC patent application no. 81 102 069.2.

The method according to the invention is a multi step method for preparing the compound of formula I and salts thereof, starting from compounds of formula II



wherein R_1 is a lower alkyl group having 1 to 6 carbon atoms, and R_2 is a lower alkyl group having 1 to 6 carbon atoms or phenyl or phenyl substituted with groups being inert under the reaction conditions of the diazotation and subsequent coupling.


Preferred lower alkyl groups are such having 1 to 3 carbon atoms such as methyl. A phenyl group R_2 can be mono- or polysubstituted by groups being inert under the reaction conditions, such as alkyl, alkoxy, halogen. Acid addition salts of the compound II can also be used as starting materials, especially salts with a mineral acid or a strong organic acid.

The starting compound of formula II is diazotized in a manner known *per se* to the corresponding diazonium salt. The diazotation is preferably carried out in conventional manner by treatment with nitrous acid in acid aqueous solution.

The formed diazonium salt of the compound II is then coupled in alkaline solution, preferably in the presence of an alkali metal hydroxide such as potassium hydroxide, with a compound of the formula III




0 036 636



III

wherein R_3 is hydrogen or a lower alkyl group having 1 to 6 carbon atoms, preferably 1 to 3 carbon atoms such as methyl.

The product formed, which corresponds to the formula IV



IV


wherein R_1 , R_2 and R_3 are as indicated above, is then isolated in a manner known *per se*, for example by acidifying the reaction mixture to precipitate the product of formula IV. It has been found that the compounds of the formula IV have excellent crystallization properties, thereby making it very simple to purify these compounds to the desired extent by recrystallization. For the recrystallization a great number of inert organic solvents can be used, for example lower aliphatic ketones such as acetone, 2-butanone, lower alkanols such as methanol, ethanol, aromatic hydrocarbons such as toluene, organic acids and esters such as acetic acid, ethyl acetate, or mixtures thereof.

The compound of formula IV, worked up to the desired purity, is then hydrolyzed in an aqueous reaction medium to the desired end product of formula I while splitting off the groups $-R_1$, $-R_3$ and $-SO_2R_2$, whereupon the end product of formula I — or salts thereof — are isolated from the reaction mixture. The hydrolysis is preferably carried out in alkaline environment, especially in the presence of an alkali metal hydroxide and possibly an inert, water-miscible solvent, for example alkanones such as acetone, alkanols such as methanol, ethanol. It would be obvious to a person skilled in the art that the hydrolysis conditions can be chosen such that the amounts of partially hydrolyzed products are minimized while at the same time no significant amounts of secondary byproducts are formed. It may, as an example, be mentioned that the hydrolysis advantageously can be carried out by boiling in water between 0.5 and 2 h using sodium hydroxide as the base. As mentioned above, and in contrast to the intermediate IV, it is comparatively difficult to purify the end product I, and it is therefore essential that the hydrolysis be carried out in the indicated manner, especially if the end product I is to be used as a drug.

After the hydrolysis has been carried out the end product is isolated from the reaction mixture in a manner known *per se*, preferably by neutralization or acidification of the reaction mixture, for example by means of a mineral acid such as hydrochloric acid or sulphuric acid, or a strong organic acid such as acetic acid.

The compound of formula I can be isolated as such or in the form of a corresponding salt, preferably a pharmaceutically acceptable salt such as an alkali metal salt. A preferred way of recovering such salts is to partially neutralize the reaction mixture with an acid to a suitable pH for precipitation of the desired salt. For example, the di-sodium and di-potassium salts can be prepared by acidifying the reaction mixture to pH 5—8, preferably 6—7, using a mineral acid or an organic acid such as acetic acid. Alternatively it is, of course, also possible to prepare the desired salts by reacting, in a manner known *per se*, the isolated acid with any suitable salt forming base.

The starting compounds of formula II used in the method according to the invention can be prepared in accordance with methods known *per se* or in analogy with methods known *per se*. For example, the compounds can be prepared by reduction of a corresponding compound of formula V



V

wherein R_1 and R_2 are as indicated above and X is a group which can be converted into an amino group, for example a nitro or arylazo group.

The invention also comprises compounds of formula IV which — as mentioned above — have proved to possess valuable crystallization properties and because thereof are excellently well suited as intermediates for the preparation of valuable end compounds such as 3,3'-zao-bis(6-hydroxy-benzoic acid) of formula I.

Although the method and the intermediates according to the invention primarily are intended for



0 036 636

preparing 3,3'-azo-bis(6-hydroxy-benzoic acid) of pharmaceutically acceptable purity, it would be obvious that they also can be used for preparing this compound for other use, wherein the requirements on purity are not as high.

The invention will now be illustrated in more detail in connection with some special non-limiting examples.

Example 1

(Preparation of starting materials)

a) Methyl 2-methane sulfonyloxy-5-nitro-benzoate

98.5 g of methyl-2-hydroxy-5-nitro-benzoate were dissolved in 250 ml of pyridine, whereupon 68.5 g of methane sulfonyl chloride were added. After heating to 50°C for 10 minutes the solution was poured on 5N ice cooled hydrochloric acid. The oil formed was extracted with chloroform. The chloroform phase was washed with water, dried and evaporated. 50 ml of toluene were added and evaporated. This was repeated once, the oil then crystallizing. Yield: 132 g of the title substance.

b) Methyl 3-amino-6-methane sulfonyloxy-benzoate.

132 g of the nitro compound from step a) were hydrogenated in 500 ml of acetic acid and 3 g of 10% palladium-on-carbon at 1—3 at. After the theoretical amount of hydrogen gas had been consumed the mixture was filtered and evaporated. It was dissolved in 1 liter of isopropanol at 65°C, whereupon HCl in ethanol was added. The hydrochloride crystallized upon cooling. After filtering the substance was shaken with sodium carbonate in water and chloroform. The chloroform phase was evaporated. The amine was re-crystallized from methanol/water. Yield: 93 g of the pure title substance.

Example 2

a) Methyl 2-hydroxy-5-[(4-methane sulfonyloxy-3-methoxycarbonylphenyl)-azo]-benzoate.

12.2 g of methyl 3-amino-6-methane sulfonyloxy-benzoate (prepared according to Example 1), 75 ml of 2N hydrochloric acid and 200 g of ice were diazotized at 0°C with 3.5 g of sodium nitrite in 20 ml of water. The diazonium salt solution was rapidly mixed with a freshly prepared solution of 15.2 g of methyl salicylate, 12 g of 85% potassium hydroxide and 1 liter of ice water while stirring vigorously. The coupling solution was acidified with hydrochloric acid after about 15 seconds. The precipitated oily product was extracted with chloroform. After drying, evaporation, and re-crystallization from 2-butanone the title substance was obtained in a yield of 12.8 g (63%). The purity was determined by liquid chromatographic analysis (HPLC) on a "reversed phase"-column (μ -Bondapak C₁₈) with methanol/water (75:25) as eluant. The total amount of impurities was only about 0.6 area percent.

b) 3,3'-azo-bis(6-hydroxy-benzoic acid).

41 g of methyl 2-hydroxy-5-[(4-methane sulfonyloxy-3-methoxycarbonylphenyl)azo]benzoate, prepared according to step a), were hydrolyzed by boiling in 800 ml of 1N sodium hydroxide for 4 h. The hydrolyzed product was precipitated by addition of concentrated hydrochloric acid to pH of about 2. The precipitate was filtered and washed with water. Yield: 29.7 g (98%) of the title substance.


Example 3

(Disodium 3,3'-azo-bis(6-hydroxy-benzoate)).

624 g of methyl 2-hydroxy-5-[(4-methane sulfonyloxy-3-methoxy carbonyl phenyl)-azo]benzoate, prepared according to step a) of Example 2, were during 15 minutes added in portions to 3 liter of water and 428 g of sodium hydroxide while boiling. After boiling for 30 minutes 360 ml of acetic acid were added during 15 minutes to pH 6. The suspension was stirred for 1.5 h without cooling, the temperature falling to 70°C. After cooling to 30°C the suspension was filtered and washed with 700 ml of methanol. After drying the title substance was obtained in a yield of 520 g (98.5%). The purity was 99.0% according to ion pair — liquid chromatographic analysis on a "reversed phase"-column (μ -Bondapak C₁₈) using methanol/water (57:43) as eluant and tetrabutyl ammonium as counter ion.

Claims


1. A method of preparing 3,3'-azo-bis-(6-hydroxy benzoic acid) of formula I




and salts thereof, comprising the steps of:
diazotizing a compound of formula II or a salt thereof




0036 636



wherein R_1 is lower alkyl and R_2 is lower alkyl, phenyl or phenyl substituted with groups being inert under the reaction conditions of the diazotation and subsequent coupling, coupling the diazonium salt obtained alkaline aqueous solution to a compound of formula III



wherein R_3 is hydrogen or lower alkyl, to form a compound of formula IV



wherein R_1 , R_2 and R_3 are as indicated above, and hydrolyzing in alkaline medium the compound of formula IV, and isolating the compound of formula I or a salt thereof.

2. The method of claim 1, wherein the coupling of the diazonium salt of the compound of formula II to the compound of formula III is carried out in the presence of an alkali metal hydroxide.

3. The method of claim 1 or 2, further comprising the step of isolating and purifying the compound of formula IV before the alkaline hydrolysis.


4. The method of claim 3, wherein the purification of the compound IV is carried out by recrystallization from an organic solvent.

5. The method of claim 1, 2, 3 or 4, wherein the hydrolysis of the compound of formula IV is carried out in the presence of an alkali metal hydroxide.

6. The method of any one of claims 1—5, comprising the step of precipitating, after the hydrolysis of the compound of formula IV, the end compound of formula I or a salt thereof by lowering the pH of the reaction mixture.

7. The method of any one of claims 1—6, for preparing the di-sodium salt of the compound of formula I, comprising the steps of hydrolyzing the compound of formula IV using sodium hydroxide and precipitating said sodium salt by neutralization to pH 5—8.

8. Compounds of formula IV




wherein R_1 is lower alkyl, R_2 is lower alkyl, phenyl or phenyl substituted with groups being inert under the conditions of a diazotation and coupling reaction and R_3 is hydrogen or lower alkyl.

9. A compound according to claim 8, wherein R_1 , R_2 and R_3 all are methyl.

10. 3,3'-azo-bis-(6-hydroxy benzoic acid) and salts thereof for use in the treatment of inflammatory intestinal diseases.

Revendications

1. Procédé de préparation d'acide 3,3'-azo-bis-(6-hydroxybenzoïque) représenté par la formule ci-après:




et de ses sels, caractérisé par les étapes suivantes:

On diazote un composé représenté par la formule (II) ci-après:



0 036 636




II

5

ou l'un de ses sels, où R_1 est un groupe alkyle inférieur et R_2 est un groupe alkyle, phényle ou phényle substitué par des groupes inertes dans les conditions de réaction de la diazotation et du couplage consécutif, on couple le sel de diazonium obtenu, dans une solution alcaline aqueuse, avec un composé représenté par la formule (III) ci-après:


15



III

dans laquelle R_3 est un atome d'hydrogène ou un radical alkyle inférieur, pour former un composé représenté par la formule (IV) ci-après:

20



IV

25

où R_1 , R_2 et R_3 sont tels que définis plus haut, l'on hydrolyse le composé de formule (IV) dans un milieu alcalin et isole le composé de formule (I) ou un sel de celui-ci.

2. Procédé selon la revendication 1, caractérisé en ce que le couplage du sel de diazonium du composé de formule (II) avec le composé de formule (III) est effectué en présence d'un hydroxyde de métal alcalin.

3. Procédé selon les revendications 1 ou 2, caractérisé en ce qu'il comprend de plus l'étape d'isolation et de purification du composé de formule (IV), avant l'hydrolyse alcaline.

4. Procédé selon la revendication 3, caractérisé en ce que la purification du composé (IV) est effectuée par recristallisation dans un solvant organique.


5. Procédé selon l'une des revendications 1 à 4, caractérisé en ce que l'hydrolyse du composé de formule (IV) est effectuée en présence d'un hydroxyde de métal alcalin.

6. Procédé selon l'une des revendications 1 à 5, caractérisé en ce qu'il comprend l'étape de précipitation, après l'hydrolyse du composé de formule (IV), du composé final de formule (I) ou d'un sel de celui-ci, par abaissement du pH du mélange de réaction.

7. Procédé selon l'une des revendications 1 à 6, caractérisé en ce que pour préparer le sel disodique du composé de formule (I), il comprend une étape d'hydrolyse du composé de formule (IV) avec de l'hydroxyde de sodium et une étape de précipitation du sel disodique par neutralisation à un pH de 5 à 8.

8. Composés caractérisés en ce qu'ils sont représentés par la formule (IV) ci-après:

50



IV

55

dans laquelle R_1 est un groupe alkyle inférieur, R_2 est un groupe alkyle inférieur phényle ou phényle substitué par des groupes inertes dans les conditions d'une réaction de diazotation et d'une réaction de couplage, et R_3 est un atome d'hydrogène ou un groupe alkyle inférieur.

9. Composé selon la revendication 8, caractérisé en ce que R_1 , R_2 et R_3 sont tous un groupe méthyle.


10. Composés selon la revendication 1, caractérisé en ce qu'il s'agit de l'acide 3,3'-azo-bis(6-hydroxybenzoïque) et de ses sels, à utiliser dans le traitement de maladies intestinales inflammatoires.

65




Patentansprüche


1. Verfahren zur Herstellung von 3,3'-Azo-bis-(6-hydroxybenzoesäure) der Formel I




und ihrer Salze, dadurch gekennzeichnet, daß man eine Verbindung der Formel II oder ein Salz davon



worin R₁ für Niedrigalkyl steht und R₂ für Niedrigalkyl, Phenyl oder Phenyl, das mit bei den Reaktionsbedingungen der Diazotisierung und nachfolgenden Kupplung inerten Gruppen substituiert ist, steht, das erhaltene Diazoniumsalz in alkalischer wäßriger Lösung zu einer Verbindung der Formel III



in der R₃ für Wasserstoff oder Niedrigalkyl steht, kuppelt, um eine Verbindung der Formel IV



worin R₁, R₂ und R₃ die obigen Bedeutungen haben, zu erhalten, und daß man die Verbindung der Formel IV im alkalischen Medium hydrolysiert und die Verbindung der Formel I oder ein Salz davon isoliert.

2. Verfahren nach Anspruch 1, dadurch gekennzeichnet, daß man das Kuppeln des Diazoniumsalzes der Verbindung der Formel II zu der Verbindung der Formel III in Gegenwart eines Alkalimetallhydroxids durchführt.

3. Verfahren nach Anspruch 1 oder 2, dadurch gekennzeichnet, daß man in einer Stufe die Verbindung der Formel IV vor der alkalischen Hydrolyse isoliert und reinigt.


4. Verfahren nach Anspruch 3, dadurch gekennzeichnet, daß man die Reinigung der Verbindung IV durch Umkristallisation aus einem organischen Lösungsmittel durchführt.

5. Verfahren nach Anspruch 1, 2, 3 oder 4, dadurch gekennzeichnet, daß man die Hydrolyse der Verbindung der Formel IV in Gegenwart eines Alkalimetallhydroxids durchführt.

6. Verfahren nach einem der Ansprüche 1 bis 5, dadurch gekennzeichnet, daß man in einer Stufe nach der Hydrolyse der Verbindung der Formel IV die Endverbindung der Formel I oder ein Salz davon durch Erniedrigung des pH-Wertes des Reaktionsgemisches ausfällt.

7. Verfahren nach einem der Ansprüche 1 bis 6, dadurch gekennzeichnet, daß man zur Herstellung des Dinatriumsalzes der Verbindung der Formel I in Stufen die Verbindung der Formel IV unter Verwendung von Natriumhydroxid hydrolysiert und das Dinatriumsalz durch Neutralisation auf einen pH-Wert von 5 bis 8 ausfällt.

8. Verbindungen der Formel IV



0 036 636

worin R_1 für Niedrigalkyl steht, R_2 für Niedrigalkyl, Phenyl oder Phenyl, das durch bei den Bedingungen der Diazotisierung und Kupplungsreaktion inerte Gruppen substituiert ist, steht und R_3 für Wasserstoff oder Niedrigalkyl steht.

- 5 9. Verbindung nach Anspruch 8, dadurch gekennzeichnet, daß R_1 , R_2 und R_3 alle Methyl sind.
10. 3,3'-Azo-bis-(6-hydroxybenzoesäure) und ihre Salze zur Verwendung bei der Behandlung von entzündlichen Darmerkrankungen.

10

15

20

25

30

35

40

45

50

55

60

65

